

Skdaccess: The **Scikit Data Access** Python Package

Quick Start Guide

v2.0.0 for Python 3.6
<https://pypi.python.org/pypi/scikit-dataaccess>


*Created and maintained by
Massachusetts Institute of Technology
Haystack Observatory, Astro- $\&$ Geo-Informatics Group*

*Project lead: Victor Pankratius
Contact email: skdaccess@mit.edu*

Code Contributors: Cody M. Rude, Justin D. Li, David M. Blair, Michael G. Gowanlock, Victor Pankratius

1 Overview

The Scikit Data Access package simplifies the handling of scientific data sets in Python. It provides a common interface across all data sets, based on a data fetcher and iterator pattern, as illustrated in the Figure below.



This paradigm places the requirements for parsing and interpreting the data inside of the data fetcher, which returns a data wrapper that provides a uniform method for accessing the data. In particular, the data wrapper implements an iterator which returns the next segment of data when requested by another function or by the user.

Advantages of Scikit Data Access

- API to import a data generator + function to get next data chunk (configurable)
- Eliminates the need to create parsers for each data set and simplifies the construction of scientific data processing pipelines.
- Enables studies involving data fusion and cross-comparisons from several sources.
- Skip parser development, dealing with physical file formats, etc.
- Can be used to download data locally or to a cloud node (e.g., Amazon Cloud). This feature simplifies distributing entire data sets or partitions of data to the cloud, and enables parallel processing in cloud computing environments.
- Easy expansion for more data sets in the future
- Code is open source (MIT License)

2 Supported Data Sets

The package introduces a common namespace and currently supports the following data sets:

Name-space	Data structure	Original Source	Data Size	Description	Download	Cache	Stream
skdaccess. astro. kepler	Dictionary of Data Frames	Mikulski Archive for Space Telescopes	$\approx 1\text{TB}$	Light curves for stars imaged by the <i>Kepler</i> Space Telescope (keplerscience.arc.nasa.gov)		x	
skdaccess. geo. groundwater	Dictionary of Data Frames	USGS National Water Information System	$\approx 1\text{GB}$	United States groundwater monitoring wells measuring the depth to water level (waterdata.usgs.gov/nwis)	x		
skdaccess. geo. pbo	Dictionary of Data Frames	UNAVCO Plate Boundary Observatory	$\approx 1\text{GB}$	Daily GPS displacement time series measurements throughout the United States (unavco.org/projects/major-projects/pbo/pbo.html)	x		
skdaccess. geo. grace	Dictionary of Data Frames	NASA Jet Propulsion Laboratory	$\approx 0.1\text{GB}$	GRACE Tellus Monthly Mass Grids. 30-day measurements of changes in Earth's gravity field to quantify equivalent water thickness (grace.jpl.nasa.gov)	x		
skdaccess. geo. gldas	Dictionary of Data Frames	NASA Jet Propulsion Laboratory	$\approx 0.1\text{GB}$	Land hydrology model produced by NASA. This version of the data is generated to match the GRACE temporal and spatial characteristics and is available as a complementary data product (grace.jpl.nasa.gov/data/get-data/land-water-content/)	x		
skdaccess. geo. modis	Dictionary of Numpy arrays	NASA MODIS	$\approx 100\text{MB}$ /image	Spectroradiometer aboard the NASA Terra and Aqua satellites that generates approximately daily images of the Earth's surface (modis.gsfc.nasa.gov)		x	x
skdaccess. geo. mahali	Dictionary of rinex files	MIT-Haystack Observatory	$\approx 10\text{GB}$	MIT led NSF project studying the Earth's ionosphere with GPS (mahali.mit.edu)		x	

3 Installation and Modes of Operation

The package can easily installed by using the standard Python “pip install” command:

```
> pip install scikit-dataaccess
```

After successful installation, a script called “skdaccess” allows users to specify the data sets that should be downloaded from their original sources to the local machine. The PBO, GRACE and groundwater data sets must be downloaded using this script before they can be used. For example, to download the PBO data use:

```
> skdaccess pbo
```

The script also completes all necessary configurations to make the data access seamlessly available in the Python environment.

3.1 Modes of Operation

There are three modes of operation available for accessing the data through the skdaccess package. The two local options are "Download" and "Cache". Using the "Download" option, the dataset is downloaded to local disk before use. The "Cache" option allows for data of interest to be downloaded during use and stored in case of future use. The online option is "Stream", which accesses the data without storing a local copy.

3.2 The Skdaccess Script

This script downloads scientific data sets from preconfigured Web sources, makes them available offline on the users machine, and configures the Python environment for data access.

For the following data sets, the skdaccess script must be used to download and prepare the data.

- GPS data from the Plate Boundary Observatory
- Depth to groundwater for wells in California
- Equivalent water thickness from GRACE Tellus Monthly Land Grids

The skdaccess script does not download Kepler data, as the data is downloaded for each star individually the first time the star is accessed by the data fetcher.

To download a dataset, use the command with the dataset name as the argument. For example, to download groundwater data available from California type

```
> skdaccess groundwater
```

The data will be downloaded into the current directory, and the .skdaccess config file located in the users home directory will be updated. Each data set can be downloaded into different directories depending on the user preferences.

To list all supported data sets, call

```
> skdaccess -l
```

This utility can install one of the following data sets:

```
PBO - Plate Boundary Observatory GPS Time Series
GRACE - Monthly Mass Grids
Groundwater - Ground water daily values from wells in California
```

Calling the script without any arguments provides a list of available commands as shown below.

```
> skdaccess
```

```
usage: skdaccess [-h] [-l] [-i LOCAL_DATA] [data_set]
```

The Sci-kit Data Access (skdaccess) package is a tool for integrating various scientific data sets into the Python environment using a

common interface. This script can download different scientific data sets for offline analysis.

positional arguments:

data_set Name of data set

optional arguments:

-h, --help show this help message and exit

-l, --list List data sets

-i LOCAL_DATA, --input LOCAL_DATA

Use LOCAL_DATA that has already been downloaded

4 Scientific Data Access in Python

Data is retrieved in a Python program via a DataFetcher object. Each data set has its own data fetcher. There are two ways of handling the data: (1) directly accessing the data structure created by the DataFetcher, or (2) through an iterator interface provided by a data wrapper.

Data Access Example:


```
# First import the data generator for water
# Note: This assumes the groundwater data has been downloaded
from skdaccess.geo.groundwater import DataFetcher as waterDF

# Create a data fetcher and get the data wrapper:
fullDF = waterDF(start_date='2007-01-01', end_date='2011-01-01')
wdata = fullDG.output().get()
```

5 Usage Examples

The following examples show how to use the data fetcher for the data sets described earlier and displaying / plotting the data. These notebooks can be accessed at <https://github.com/MITHaystack/scikit-dataaccess/tree/master/skdaccess/examples>.

5.1 skdaccess.astro.kepler



5.2 skdaccess.geo.groundwater

MIT Computer-Aided Discovery Demo_Groundwater Last Checkpoint: 5 hours ago (autosaved) Control Panel Logout

File Edit View Insert Cell Kernel Help Python 3

Code Cell Toolbar: None

```
In [1]: %matplotlib notebook
import matplotlib.pyplot as plt

In [2]: # USGS Groundwater Data - 129 Monitoring Wells in CA between 2010 and 2014
# Source: http://water.usgs.gov/ogw/
# Returns depth to water level in meters

In [3]: from skdaccess.geo.groundwater import DataFetcher as GW_DF

In [4]: groundwater_fetcher = GW_DF(start_date='2010-01-01',end_date='2014-01-01')

In [5]: groundwater_data = groundwater_fetcher.output().get() # returns a pandas data panel

In [6]: groundwater_data.loc['Water Depth'].head()
```

	323313117033901	323313117033902	323313117033903	323313117033904	323313117033905	323527117050001	323527117050002	323
2010-01-01	12.896088	12.981432	13.024104	5.830824	NaN	13.203936	10.250424	6.14
2010-01-02	12.877800	12.960096	13.005816	5.797296	NaN	13.200888	10.146792	6.08
2010-01-03	12.877800	12.960096	13.002768	5.775960	NaN	13.213080	10.064496	6.05
2010-01-04	12.877800	12.960096	13.005816	5.760720	NaN	13.219176	9.982200	6.01
2010-01-05	12.877800	12.960096	13.005816	5.739384	NaN	13.222224	9.909048	5.97


5 rows x 129 columns

```
In [7]: # Plotting Well Number 323313117033902
plt.figure();
plt.plot(groundwater_data.loc['Water Depth'],, '323313117033902');
plt.tight_layout()
```


Figure 1

x=Jul 2010 y=12.7499

5.3 skdaccess.geo.pbo



5.4 skdaccess.geo.grace



Acknowledgements

Many thanks for support from NASA AIST NNX15AG84G, NSF ACI 1442997, NSF AGS-1343967, and the Amazon Web Services Research grants (PI: V. Pankratius).